## MUC1(CT) Antibody

Catalog No: #43222



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _  |    |     |     |              |   |
|----|----|-----|-----|--------------|---|
| רו | es | Cri | nt  | $\mathbf{a}$ | n |
| u  | -5 | C I | IVI | II U         |   |

| Product Name          | MUC1(CT) Antibody   |
|-----------------------|---|
| Host Species          | Rabbit  |
| Clonality             | Polyclonal  |
| Purification          | Antigen affinity purification.  |
| Applications          | IHC   |
| Species Reactivity    | Hu  |
| Specificity           | The antibody detects endogenous levels of total MUC1(CT) protein.               |
| Immunogen Type        | peptide   |
| Immunogen Description | Synthetic peptide of human MUC1(CT)   |
| Target Name           | MUC1(CT)  |
| Other Names           | EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; |
|                       | ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC                                   |
| Accession No.         | Swiss-Prot#: P15941?Gene ID: 4582   |
| Uniprot               | P15941  |
| GeneID                | 4582;   |
| Concentration         | 1.2mg/ml  |
| Formulation           | Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.                              |
| Storage               | Store at -20°C  |
|                       |   |


## Application Details

Immunohistochemistry: 1:25-1:100

## **Images**



Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue using #43222 at dilution 1/20.



Immunohistochemical analysis of paraffin-embedded Human esophagus cancer tissue using #43222 at dilution 1/20.

## Background

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.

Note: This product is for in vitro research use only