CCL7 Antibody

Catalog No: #43233



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_			
	escri	ınt	ion.
\boldsymbol{L}	COUL	Iυι	IUII

CCL7 Antibody
Rabbit
Polyclonal
Antigen affinity purification.
IHC
Hu
The antibody detects endogenous levels of total CCL7 protein.
peptide
Synthetic peptide of human CCL7
CCL7
FIC; MARC; MCP3; NC28; MCP-3; SCYA6; SCYA7
Swiss-Prot#: P80098Gene ID: 6354
P80098
6354;
0.8mg/ml
Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
Store at -20°C

Application Details

Immunohistochemistry: 1:100-1:200

Images



Immunohistochemical analysis of paraffin-embedded Human Liver cancer tissue using #43233 at dilution 1/200.

Background

This gene encodes monocyte chemotactic protein 3, a secreted chemokine which attracts macrophages during inflammation and metastasis. It is a member of the C-C subfamily of chemokines which are characterized by having two adjacent cysteine residues. The protein is an in vivo substrate of matrix metalloproteinase 2, an enzyme which degrades components of the extracellular matrix. This gene is part of a cluster of C-C chemokine family members on chromosome 17q.

Note: This product is for in vitro research use only