

# Calcium Sensing Receptor (Phospho-Thr888) Conjugated Antibody



Catalog No: #C12041

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

Package Size: #C12041-AF350 100ul #C12041-AF405 100ul #C12041-AF488 100ul

#C12041-AF555 100ul #C12041-AF594 100ul #C12041-AF647 100ul

#C12041-AF680 100ul #C12041-AF750 100ul #C12041-Biotin 100ul

## Description

Product Name	Calcium Sensing Receptor (Phospho-Thr888) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of Calcium Sensing Receptor only when phosphorylated at Threonine 888.
Immunogen Description	Peptide sequence around phosphorylation site of Threonine 888 (R-A-T(p)-L-R) derived from Human Calcium Sensing Receptor.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	CAR;FHH;FIH;HHC;EIG8
Accession No.	Swiss-Prot#:P41180NCBI Gene ID:846NCBI mRNA#:NM_000388.3NCBI Protein#:NP_000379.2
Uniprot	P41180
GeneID	846;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	140
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

---

## Product Description

---

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

## Background

---

Senses changes in the extracellular concentration of calcium ions. The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system.

---

Note: This product is for in vitro research use only