

## RAD17 Conjugated Antibody

Catalog No: #C42728



Package Size: #C42728-AF350 100ul #C42728-AF405 100ul #C42728-AF488 100ul  
 #C42728-AF555 100ul #C42728-AF594 100ul #C42728-AF647 100ul  
 #C42728-AF680 100ul #C42728-AF750 100ul #C42728-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

|                       |  |
|-----------------------|--|
| Product Name          | RAD17 Conjugated Antibody  |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Species Reactivity    | Hu   |
| Specificity           | The antibody detects endogenous levels of total RAD17 protein.   |
| Immunogen Description | Fusion protein of human RAD17  |
| Conjugates            | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750   |
| Other Names           | CCYC; R24L; RAD24; HRAD17; RAD17SP   |
| Accession No.         | Swiss-Prot#:O75943NCBI Gene ID:5884NCBI mRNA#:BC032304NCBI Protein#:   |
| Uniprot               | O75943   |
| GeneID                | 5884;  |
| Excitation Emission   | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Calculated MW         | 77KD   |
| Formulation           | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide  |
| Storage               | Store at 4°C in dark for 6 months  |

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

The protein encoded by this gene is highly similar to the gene product of *Schizosaccharomyces pombe rad17*, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This protein shares strong similarity with DNA replication factor C (RFC), and can form a complex with RFCs. This protein binds to chromatin prior to DNA damage and is phosphorylated by the checkpoint kinase ATR following damage.

---

Note: This product is for in vitro research use only