ATP1B2 Conjugated Antibody

Catalog No: #C43972



 Package Size:
 #C43972-AF350 100ul
 #C43972-AF405 100ul
 #C43972-AF488 100ul

 #C43972-AF555 100ul
 #C43972-AF594 100ul
 #C43972-AF647 100ul

 #C43972-AF680 100ul
 #C43972-AF750 100ul
 #C43972-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	ATP1B2 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous levels of total ATP1B2 protein.
Immunogen Description	Synthetic peptide of human ATP1B2
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	AMOG
Accession No.	Swiss-Prot#:P14415NCBI Gene ID:482NCBI mRNA#:NCBI Protein#:NP_001669
Uniprot	P14415
GeneID	482;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	33
Formulation	Liquid in PBS containing 40% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at 4°Cin dark for 6 months

Application Details

Suggested Dilution:	
AF350 conjugated: most applications: 1: 50 - 1: 250	
AF405 conjugated: most applications: 1: 50 - 1: 250	
AF488 conjugated: most applications: 1: 50 - 1: 250	
AF555 conjugated: most applications: 1: 50 - 1: 250	
AF594 conjugated: most applications: 1: 50 - 1: 250	
AF647 conjugated: most applications: 1: 50 - 1: 250	
AF680 conjugated: most applications: 1: 50 - 1: 250	
AF750 conjugated: most applications: 1: 50 - 1: 250	
Piotin conjugated: working with onzyme conjugated strontavidin, most applications: 1:50, 1:1,000	

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The protein encoded by this gene belongs to the family of Na+/K+ and H+/K+ ATPases beta chain proteins, and to the subfamily of Na+/K+ -ATPases. Na+/K+ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane. The glycoprotein subunit of Na+/K+ -ATPase is encoded by multiple genes. This gene encodes a beta 2 subunit. Two transcript variants encoding different isoforms have been found for this gene.

Note: This product is for in vitro research use only