ANKRD27 Antibody

Catalog No: #43897



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


110	10 Cri	Intion	
	וטסי	iption	
_			

ANKRD27 Antibody	
Rabbit	
Polyclonal	
Antigen affinity purification	
IHC	
Hu	
The antibody detects endogenous levels of total ANKRD27 protein.	
protein	
Fusion protein of human ANKRD27	
ANKRD27	
VARP; PP12899	
Swiss-Prot#: Q96NW4NCBI Gene ID: 84079	
Q96NW4	
84079;	
0.5mg/ml	
Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.	
Store at -20°C	

Application Details

Immunohistochemistry: 1: 20-100

Images



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ANKRD27 Antibody at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x200)

Background

May be a guanine exchange factor (GEF) for Rab21, Rab32 and Rab38 and regulate endosome dynamics (PubMed:16525121, PubMed:18477474). May regulate the participation of VAMP7 in membrane fusion events; in vitro inhibits VAMP7-mediated SNARE complex formation by trapping VAMP7 in a closed, fusogenically inactive conformation (PubMed:23104059). Involved in peripheral melanosomal distribution of TYRP1 in melanocytes; the function, which probably is implicating vesicle-trafficking, includes cooperation with Rab32, Rab38 and VAMP7 (By similarity). Involved in the

regulation of neurite growth; the function seems to require its GEF activity, probably towards Rab21, and VAMP7 but not Rab32/38 (By similarity). Proposed to be involved in Golgi sorting of VAMP7 and transport of VAMP7 vesicles to the cell surface; the function seems to implicate kinesin heavy chain isoform 5 proteins, GOLGA4, RAB21 and MACF1 (PubMed:22705394). Required for the colocalization of VAMP7 and Rab21, probably on TGN sites (PubMed:19745841). Involved in GLUT1 endosome-to-plasma membrane trafficking; the function is dependent of association with VPS29 (PubMed:24856514).

Note: This product is for in vitro research use only