

## BDH2 Conjugated Antibody

Catalog No: #C46344



Package Size: #C46344-AF350 100ul #C46344-AF405 100ul #C46344-AF488 100ul  
 #C46344-AF555 100ul #C46344-AF594 100ul #C46344-AF647 100ul  
 #C46344-AF680 100ul #C46344-AF750 100ul #C46344-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

|                       |  |
|-----------------------|--|
| Product Name          | BDH2 Conjugated Antibody   |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Species Reactivity    | Hu Ms  |
| Specificity           | The antibody detects endogenous levels of total BDH2 protein.  |
| Immunogen Description | Full length fusion protein of human BDH2   |
| Conjugates            | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750   |
| Other Names           | DHRS6; EFA6R; SDR15C1; UCPA-OR; UNQ6308; PRO20933  |
| Accession No.         | Swiss-Prot#:Q9BUT1NCBI Gene ID:56898NCBI mRNA#:NCBI Protein#:BC001953  |
| Uniprot               | Q9BUT1   |
| GeneID                | 56898;   |
| Excitation Emission   | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Calculated MW         | 27   |
| Formulation           | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide  |
| Storage               | Store at 4°C in dark for 6 months  |

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

Dehydrogenase that mediates the formation of 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin and associates with LCN2, thereby playing a key role in iron homeostasis and transport. Also acts as a 3-hydroxybutyrate dehydrogenase (By similarity).

---

Note: This product is for in vitro research use only