CPM Conjugated Antibody

Catalog No: #C46545

SAB Signalway Antibody

Package Size: #C46545-AF350 100ul #C46545-AF405 100ul #C46545-AF488 100ul

#C46545-AF555 100ul #C46545-AF594 100ul #C46545-AF647 100ul

#C46545-AF680 100ul #C46545-AF750 100ul #C46545-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | CPM Conjugated Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total CPM protein. |
| Immunogen Description | Synthetic protein corresponding to residues near the C terminal of human CPM |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Accession No. | Swiss-Prot#:P14384NCBI Gene ID:1368NCBI mRNA#:NCBI Protein#:BC022276 |
| Uniprot | P14384 |
| GeneID | 1368; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 51 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°Cin dark for 6 months |
| | |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

 $Biotin \ conjugated: working \ with \ enzyme-conjugated \ streptavidin, \ most \ applications: \ 1:50 - 1:1,000$

Background

The protein encoded by this gene is a membrane-bound arginine/lysine carboxypeptidase. Its expression is associated with monocyte to macrophage differentiation. This encoded protein contains hydrophobic regions at the amino and carboxy termini and has 6 potential asparagine-linked glycosylation sites. The active site residues of carboxypeptidases A and B are conserved in this protein. Three alternatively spliced transcript variants encoding the same protein have been described for this gene.

Note: This product is for in vitro research use only