

XRCC4 antibody

Catalog No: #22951

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	XRCC4 antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC IF
Species Reactivity	Hu
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide contain a sequence corresponding to a region within amino acids 216 and 269 (Q13426) of XRCC4
Target Name	XRCC4
Accession No.	Swiss-Prot:Q13426Gene ID:7518
Uniprot	Q13426
GeneID	7518;
Concentration	1mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 38kd

Western blotting: 1:500-1:3000


Immunohistochemistry: 1:100-1:250

Immunofluorescence: 1:100-1:200


Images



Sample(30 ug whole cell lysate)
 A: 293T
 B: A431
 C: HeLa S3
 10% SDS PAGE
 Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded A549 xenograft, using XRCC4 antibody at 1: 100 dilution.



Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using XRCC4 antibody at 1: 200 dilution.

Background

The protein encoded by this gene functions together with DNA ligase IV and the DNA-dependent protein kinase in the repair of DNA double-strand break by non-homologous end joining and the completion of V(D)J recombination events. The non-homologous end-joining pathway is required both for normal development and for suppression of tumors. This gene functionally complements XR-1 Chinese hamster ovary cell mutant, which is impaired in DNA double-strand breaks produced by ionizing radiation and restriction enzymes. Alternative transcription initiation and alternative splicing generates several transcript variants. [provided by RefSeq]

Note: This product is for in vitro research use only