

HER2 (Phospho-Tyr1221/Tyr1222) Conjugated Antibody

Catalog No: #C11076

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C11076-AF350 100ul #C11076-AF405 100ul #C11076-AF488 100ul

#C11076-AF555 100ul #C11076-AF594 100ul #C11076-AF647 100ul

#C11076-AF680 100ul #C11076-AF750 100ul #C11076-Biotin 100ul

Description

Product Name	HER2 (Phospho-Tyr1221/Tyr1222) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of HER2 only when phosphorylated at tyrosine1221/tyrosine1222.
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine1221/1222 (N-L-Y(p)-Y(p)-W) derived from Human HER2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	C-erbB-2;ErbB2
Accession No.	Swiss-Prot#:P04626NCBI Gene ID:2064NCBI mRNA#:NM_001005862.1 NCBI Protein#: NP_001005862.1
Uniprot	P04626
GeneID	2064;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	185
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Not activated by EGF, TGF-alpha and amphiregulin.

Note: This product is for in vitro research use only