

NMDAR1 (Phospho-Ser896) Conjugated Antibody

Catalog No: #C11104



Package Size: #C11104-AF350 100ul #C11104-AF405 100ul #C11104-AF488 100ul
 #C11104-AF555 100ul #C11104-AF594 100ul #C11104-AF647 100ul
 #C11104-AF680 100ul #C11104-AF750 100ul #C11104-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	NMDAR1 (Phospho-Ser896) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of NMDAR1 only when phosphorylated at serine896.
Immunogen Description	Peptide sequence around phosphorylation site of serine 896 (R-R-S(p)-S-K) derived from Human NMDAR1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	GLURZ1;GRIN1;NMD-R1;NMDZ1;NMZ1
Accession No.	Swiss-Prot#:Q05586NCBI Gene ID:2902NCBI mRNA#:NM_000832.5NCBI Protein#:NP_000823.4
Uniprot	Q05586
GeneID	2902;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	120
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

NMDA receptors are members of the ionotropic class of glutamate receptors, which also includes Kainate and AMPA receptors. NMDA receptors consist of NR1 subunits combined with one or more NR2 (A-D) or NR3 (A-B) subunits. The ligand-gated channel is permeable to cations including Ca^{2+} , and at resting membrane potentials NMDA receptors are inactive due to a voltage-dependent blockade of the channel pore by Mg^{2+} . NMDA receptor activation, which requires binding of glutamate and glycine, leads to an influx of Ca^{2+} into the postsynaptic region where it activates several signaling cascades, including pathways leading to the induction of long-term potentiation (LTP) and depression (LTD). NMDA receptors have a critical role in excitatory synaptic transmission and plasticity in the CNS. They govern a range of physiological conditions including neurological disorders caused by excitotoxic neuronal injury, psychiatric disorders and neuropathic pain syndromes.

Note: This product is for in vitro research use only