

Synaptotagmin 1/2 (Phospho-Thr202/199) Conjugated Antibody

Catalog No: #C11210

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C11210-AF350 100ul #C11210-AF405 100ul #C11210-AF488 100ul

#C11210-AF555 100ul #C11210-AF594 100ul #C11210-AF647 100ul

#C11210-AF680 100ul #C11210-AF750 100ul #C11210-Biotin 100ul

Description

Product Name	Synaptotagmin 1/2 (Phospho-Thr202/199) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of Synaptotagmin 1/2 only when phosphorylated at threonine202/199.
Immunogen Description	Peptide sequence around phosphorylation site of threonine 202/199 (R-K-T(p)-L-N) derived from Human Synaptotagmin 1/2 .
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	SYT1/2; SytI/II; Synaptotagmin I/II; P65 ; SYT
Accession No.	Swiss-Prot#:P21579/Q8N9I0NCBI Gene ID:6857/127833NCBI mRNA#:NM_001135805.1 NM_001136504.1NCBI Protein#: NP_001129277.1 NP_001129976.1
Uniprot	P21579
GeneID	6857;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	46
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

The synaptotagmins are integral membrane proteins of synaptic vesicles thought to serve as Ca^{2+} sensors in the process of vesicular trafficking and exocytosis. Calcium binding to synaptotagmin I participates in triggering neurotransmitter release at the synapse

Note: This product is for in vitro research use only