Product Datasheet

p44/42 MAP Kinase (Phospho-Thr202) Conjugated Antibody

Catalog No: #C11245



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size:	#C11245-AF350 100ul	#C11245-AF405 100ul	#C11245-AF488 100ul
	#C11245-AF555 100ul	#C11245-AF594 100ul	#C11245-AF647 100ul
	#C11245-AF680 100ul	#C11245-AF750 100ul	#C11245-Biotin 100ul

Description

Product Name	p44/42 MAP Kinase (Phospho-Thr202) Conjugated Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Species Reactivity	Hu Ms Rt		
Specificity	The antibody detects endogenous level of p44/42 MAP Kinase only when phosphorylated at threonine 202.		
Immunogen Description	Peptide sequence around phosphorylation site of threonine 202 (F-L-T(p)-E-Y) derived from Human p44/42		
	MAP Kinase.		
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750		
Other Names	ERK;ERT2;Extracellular signal- regulated kinase 1;Insulin-stimulated MAP2 kinase;MAP kinase 1		
Accession No.	Swiss-Prot#:P27361NCBI Gene ID:5595NCBI mRNA#:NM_001040056.1 NCBI Protein#:NP_001035145.1		
Uniprot	P27361		
GeneID	5595;		
Excitation Emission	AF350: 346nm/442nm		
	AF405: 401nm/421nm		
	AF488: 493nm/519nm		
	AF555: 555nm/565nm		
	AF594: 591nm/614nm		
	AF647: 651nm/667nm		
	AF680: 679nm/702nm		
	AF750: 749nm/775nm		
Calculated MW	42 44		
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide		
Storage	Store at 4°C in dark for 6 months		

Application Details

Suggested Dilution:			
AF350 conjugated: most applications: 1: 50 - 1: 250			
AF405 conjugated: most applications: 1: 50 - 1: 250			
AF488 conjugated: most applications: 1: 50 - 1: 250			
AF555 conjugated: most applications: 1: 50 - 1: 250			
AF594 conjugated: most applications: 1: 50 - 1: 250			
AF647 conjugated: most applications: 1: 50 - 1: 250			
AF680 conjugated: most applications: 1: 50 - 1: 250			

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1.

Note: This product is for in vitro research use only