

α -Synuclein (Phospho-Tyr125) Conjugated Antibody

Catalog No: #C11256



Package Size: #C11256-AF350 100ul #C11256-AF405 100ul #C11256-AF488 100ul
 #C11256-AF555 100ul #C11256-AF594 100ul #C11256-AF647 100ul
 #C11256-AF680 100ul #C11256-AF750 100ul #C11256-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	α -Synuclein (Phospho-Tyr125) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of α -Synuclein only when phosphorylated at tyrosine 125.
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 125 (E-A-Y(p)-E-M) derived from Human α -Synuclein.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	NACP;SYN;SYUA;alpha-synuclein
Accession No.	Swiss-Prot#:P37840NCBI Gene ID:6622NCBI mRNA#:NM_000345.3NCBI Protein#:NP_000336.1
Uniprot	P37840
GeneID	6622;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	18
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

SncA is a member of the synuclein family of structurally related proteins that are prominently expressed in the central nervous system, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and SncA and Snc-Beta inhibit phospholipase D2 selectively. SncA may serve to integrate presynaptic signaling and membrane trafficking. Aggregated SncA proteins form brain lesions that are hallmarks of neurodegenerative synucleinopathies. Defects in SncA play a role in the pathogenesis of Parkinson disease. SncA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer disease. SncA shares 95% sequence homology with rat SncA. Rat SncA is specifically expressed in brain and is associated with synaptosomal membranes in neurons.

Note: This product is for in vitro research use only