

SHP-2 (Phospho-Tyr580) Conjugated Antibody

Catalog No: #C11320



Package Size: #C11320-AF350 100ul #C11320-AF405 100ul #C11320-AF488 100ul
 #C11320-AF555 100ul #C11320-AF594 100ul #C11320-AF647 100ul
 #C11320-AF680 100ul #C11320-AF750 100ul #C11320-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	SHP-2 (Phospho-Tyr580) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of SHP-2 only when phosphorylated at tyrosine 580.
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 580 (R-V-Y(p)-E-N) derived from Human SHP-2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	PTN11;PTP-1D;PTP-2C;PTP2C;PTPN11
Accession No.	Swiss-Prot#:Q06124NCBI Gene ID:5781NCBI mRNA#:NM_002834.3 NCBI Protein#:NP_002825.3
Uniprot	Q06124
GeneID	5781;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	72
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus.

Note: This product is for in vitro research use only