

Acetyl-CoA Carboxylase (Phospho-Ser79) Conjugated Antibody



Catalog No: #C11584

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C11584-AF350 100ul #C11584-AF405 100ul #C11584-AF488 100ul

#C11584-AF555 100ul #C11584-AF594 100ul #C11584-AF647 100ul

#C11584-AF680 100ul #C11584-AF750 100ul #C11584-Biotin 100ul

Description

Product Name	Acetyl-CoA Carboxylase (Phospho-Ser79) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of Acetyl-CoA Carboxylase only when phosphorylated at serine 79.
Immunogen Description	Peptide sequence around phosphorylation site of serine 79(S-M-S(p)-G-L) derived from Human Acetyl-CoA Carboxylase.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	ACC;ACAC;ACC1;ACCA
Accession No.	Swiss-Prot#:Q13085NCBI Gene ID:31NCBI mRNA#:NM_198834.1NCBI Protein#:NP_942131.1
Uniprot	Q13085
GeneID	31;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	265
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Catalyzes the rate-limiting reaction in the biogenesis of long-chain fatty acids. Carries out three functions: biotin carboxyl carrier protein, biotin carboxylase and carboxyltransferase.

Note: This product is for in vitro research use only