

p47 phox (Phospho-Ser345) Conjugated Antibody

Catalog No: #C11811



Package Size: #C11811-AF350 100ul #C11811-AF405 100ul #C11811-AF488 100ul
 #C11811-AF555 100ul #C11811-AF594 100ul #C11811-AF647 100ul
 #C11811-AF680 100ul #C11811-AF750 100ul #C11811-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	p47 phox (Phospho-Ser345) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of p47 phox only when phosphorylated at serine 345.
Immunogen Description	Peptide sequence around phosphorylation site of Serine 345(P-Q-S(p)-P-G) derived from Human Nuclear p47 phox.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	NCF1; P47 phox;NCF-47K
Accession No.	Swiss-Prot#:P14598NCBI Gene ID:653361NCBI mRNA#:NM_000265.5. NCBI Protein#:NP_000256.4.
Uniprot	P14598
GeneID	653361;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	45
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production).

Note: This product is for in vitro research use only