

Smad2 (Phospho-Ser465) Conjugated Antibody

Catalog No: #C11838



Package Size: #C11838-AF350 100ul #C11838-AF405 100ul #C11838-AF488 100ul
 #C11838-AF555 100ul #C11838-AF594 100ul #C11838-AF647 100ul
 #C11838-AF680 100ul #C11838-AF750 100ul #C11838-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	Smad2 (Phospho-Ser465) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of Smad2 only when phosphorylated at serine 465.
Immunogen Description	Peptide sequence around phosphorylation site of Serine 465(C-S-S(p)-M-S) derived from Human Smad2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	hMAD-2;JV18;JV18-1; MAD;hSMAD2
Accession No.	Swiss-Prot#:Q15796NCBI Gene ID:4087NCBI mRNA#:NM_001003652.3. NCBI Protein#:NP_001003652.1.
Uniprot	Q15796
GeneID	4087;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	52
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Transcriptional modulator activated by TGF-beta and activin type 1 receptor kinase. SMAD2 is a receptor-regulated SMAD (R-SMAD). May act as a tumor suppressor in colorectal carcinoma.

Note: This product is for in vitro research use only