

## MNK1 (Phospho-Thr255) Conjugated Antibody

Catalog No: #C12142



Package Size: #C12142-AF350 100ul #C12142-AF405 100ul #C12142-AF488 100ul  
 #C12142-AF555 100ul #C12142-AF594 100ul #C12142-AF647 100ul  
 #C12142-AF680 100ul #C12142-AF750 100ul #C12142-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	MNK1 (Phospho-Thr255) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of MNK1 only when phosphorylated at threonine 255.
Immunogen Description	Peptide sequence around phosphorylation site of threonine 255 (L-T-T(p)-P-C) derived from Human MNK1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	EC 2.7.11.1; kinase Mnk1; Map kinase interacting kinase; MAP kinase signal-integrating kinase 1; MAP kinase-interacting serine/threonine kinase 1; MKNK1
Accession No.	Swiss-Prot#:Q9BUB5NCBI Gene ID:8569
Uniprot	Q9BUB5
GeneID	8569;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	60
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250  
 AF405 conjugated: most applications: 1: 50 - 1: 250  
 AF488 conjugated: most applications: 1: 50 - 1: 250  
 AF555 conjugated: most applications: 1: 50 - 1: 250  
 AF594 conjugated: most applications: 1: 50 - 1: 250  
 AF647 conjugated: most applications: 1: 50 - 1: 250  
 AF680 conjugated: most applications: 1: 50 - 1: 250  
 AF750 conjugated: most applications: 1: 50 - 1: 250

## Product Description

---

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

## Background

---

May play a role in the response to environmental stress and cytokines. Appears to regulate translation by phosphorylating EIF4E, thus increasing the affinity of this protein for the 7-methylguanosine-containing mRNA cap.

---

Note: This product is for in vitro research use only