

Histone H3(Phospho-S10) Conjugated Antibody

Catalog No: #C13337



Package Size: #C13337-AF350 100ul #C13337-AF405 100ul #C13337-AF488 100ul
 #C13337-AF555 100ul #C13337-AF594 100ul #C13337-AF647 100ul
 #C13337-AF680 100ul #C13337-AF750 100ul #C13337-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	Histone H3(Phospho-S10) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	p53 like transcription factor antibody p53 related protein antibody p53-like transcription factor antibody p53-related protein antibody p73 antibody P73_HUMAN antibody TP73 antibody Tumor protein p73 antibody
Accession No.	Swiss-Prot#:O15350
Uniprot	O15350
GeneID	7161;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	70/60 (12 isoforms)
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The p53 gene is a widely studied anti-oncogene, or tumor suppressor gene. The p53 gene product can act as a negative regulator of cell growth in response to DNA damage. Mutations and allelic loss of the p53 gene have been associated with malignant transformation in a wide variety of human tumors. p53 shares considerable sequence similarity with p73, a gene that maps to a region in chromosome 1 that is frequently deleted in neuroblastomas. However, p73 does not appear to be activated by DNA damaging agents. The p73 isoform p73 α inhibits drug-induced apoptosis in small cell lung carcinoma cells, while the p73 isoform p73 β promotes it. p73 α also prevents Bax activation, mitochondrial dysfunction, caspase activation and is able to reduce apoptosis induced by the BH3-only protein PUMA (p53 upregulated modulator of apoptosis). There is an equilibrium between p73 α and p73 β , demonstrated by the fact that p73 α inhibits the pro-apoptotic effect of p73 β .

Note: This product is for in vitro research use only