

MEK1(Phospho-S218/S222) Conjugated Antibody

Catalog No: #C13373



Package Size: #C13373-AF350 100ul #C13373-AF405 100ul #C13373-AF488 100ul
 #C13373-AF555 100ul #C13373-AF594 100ul #C13373-AF647 100ul
 #C13373-AF680 100ul #C13373-AF750 100ul #C13373-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	MEK1(Phospho-S218/S222) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	I kappa B alpha antibody I-kappa-B-alpha antibody IkappaBalpha antibody IkB-alpha antibody IKBA antibody IKBA_HUMAN antibody IKBalpha antibody MAD 3 antibody MAD3 antibody Major histocompatibility complex enhancer-binding protein MAD3 antibody NF kappa B inhibitor alpha antibody NF-kappa-B inhibitor alpha antibody NFKBI antibody NFKBIA antibody Nuclear factor of kappa light chain gene enhancer in B cells antibody Nuclear factor of kappa light polypeptide gene enhancer in B cells inhibitor alpha antibody
Accession No.	Swiss-Prot#:P25963
Uniprot	P25963
GeneID	4792;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	34
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins, which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime.

Note: This product is for in vitro research use only