

## Hsp27(Phospho-S82) Conjugated Antibody

Catalog No: #C13388



Package Size: #C13388-AF350 100ul #C13388-AF405 100ul #C13388-AF488 100ul  
 #C13388-AF555 100ul #C13388-AF594 100ul #C13388-AF647 100ul  
 #C13388-AF680 100ul #C13388-AF750 100ul #C13388-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Hsp27(Phospho-S82) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	CDC 37 antibody Cdc37 antibody CDC37 cell division cycle 37 homolog antibody CDC37 cell division cycle 37 S cerevisiae homolog antibody CDC37 cell division cycle 37, S cerevisiae, homolog of antibody Cdc37 homolog antibody CDC37 protein antibody CDC37_HUMAN antibody CDC37A antibody cell division cycle 37 antibody Cell division cycle 37 homolog antibody Hsp90 chaperone protein kinase targeting subunit antibody Hsp90 chaperone protein kinase targeting subunit p50Cdc37 antibody Hsp90 chaperone protein kinase-targeting subunit antibody Hsp90 co chaperone Cdc37 antibody Hsp90 co-chaperone Cdc37 antibody p50 antibody p50Cdc37 antibody S cerevisiae hypothetical protein CDC37 antibody
Accession No.	Swiss-Prot#:Q16543
Uniprot	Q16543
GeneID	11140;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	45
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

---

## Background

Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by the proteolysis of cyclins. The cell division cycle (Cdc) genes are required at various points in the cell cycle. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. Cdc6 is the human homolog of *Saccharomyces cerevisiae* Cdc6, which is involved in the initiation of DNA replication. Cdc37 appears to facilitate Cdk4/cyclin D1 complex formation and has been shown to form a stable complex with Hsp90. Cdc34, Cdc27 and Cdc16 function as ubiquitin-conjugating enzymes. Cdc34 is thought to be the structural and functional homolog of *Saccharomyces cerevisiae* Cdc34, which is essential for the G1 to S phase transition. Cdc16 and Cdc27 are components of the APC (anaphase-promoting complex) which ubiquitinates cyclin B, resulting in cyclin B/Cdk complex degradation.

---

Note: This product is for in vitro research use only