

EGFR(Phospho-S695) Conjugated Antibody

Catalog No: #C13398



Package Size: #C13398-AF350 100ul #C13398-AF405 100ul #C13398-AF488 100ul
 #C13398-AF555 100ul #C13398-AF594 100ul #C13398-AF647 100ul
 #C13398-AF680 100ul #C13398-AF750 100ul #C13398-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	EGFR(Phospho-S695) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Mammary gland factor (STAT5A) antibody MGF (STAT5A) antibody Signal transducer and activator of transcription 5A antibody Signal transducer and activator of transcription 5B antibody STA5A_HUMAN antibody Stat 5a antibody STAT 5b antibody STAT5A antibody Transcription factor STAT5B antibody
Accession No.	Swiss-Prot#:P42229
Uniprot	P42229
GeneID	6776;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	90
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Background

Stat5 (Signal Transducers and Activators of Transcription 5) is important in regulating T cell functions involving the receptors for Interleukin-2 (IL-2). IL-2 stimulates the rapid phosphorylation of both serine and tyrosine residues of Stat5a and Stat5b in human T lymphocytes and in several IL-2-responsive lymphocytic cell lines. IL-2 differentially induces serine phosphorylation of Stat5a and Stat5b on Ser726 and Ser731, respectively. Stat5b is preferentially phosphorylated and displays more protracted serine phosphorylation kinetics than Stat5a. Both the acid-rich region and the COOH terminus of IL-2R β can independently mediate IL-2-induced Stat 5a/b serine phosphorylation, suggesting that Stat5a/b serine phosphorylation occurs at a postreceptor level. Stat5a is phosphorylated on Tyr694 in a prolactin-sensitive manner, whereas serine phosphorylation is constitutive. Activation of Stat5 by IL-2 may help govern the biological effects of IL-2 during the immune response.

Note: This product is for in vitro research use only