

## MEK-2 (Ab-394) Conjugated Antibody

Catalog No: #C21008



Package Size: #C21008-AF350 100ul #C21008-AF405 100ul #C21008-AF488 100ul  
 #C21008-AF555 100ul #C21008-AF594 100ul #C21008-AF647 100ul  
 #C21008-AF680 100ul #C21008-AF750 100ul #C21008-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	MEK-2 (Ab-394) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Rt
Specificity	The antibody detects endogenous level of total MEK2 protein.
Immunogen Description	Peptide sequence around aa.392~396 (P-G-T-P-T) derived from Human MEK-2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	ERK activator kinase 2;MAP kinase kinase 2;MAP2K2;MAPK/ERK kinase 2;MAPKK 2
Accession No.	Swiss-Prot#:P36507NCBI Gene ID:5605NCBI mRNA#:NM_030662.3 NCBI Protein#:NP_109587.1
Uniprot	P36507
GeneID	5605;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	44
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Product Description

---

Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.

## Background

---

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases

---

Note: This product is for in vitro research use only