

PAK1/PAK2/PAK3 (Ab-423/402/421) Conjugated Antibody

Catalog No: #C21169

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C21169-AF350 100ul #C21169-AF405 100ul #C21169-AF488 100ul

#C21169-AF555 100ul #C21169-AF594 100ul #C21169-AF647 100ul

#C21169-AF680 100ul #C21169-AF750 100ul #C21169-Biotin 100ul

Description

Product Name	PAK1/PAK2/PAK3 (Ab-423/402/421) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total PAK1/PAK2/PAK3 protein.
Immunogen Description	Peptide sequence around aa.421~425/400~404/419~423 (R-S-T-M-V) derived from Human PAK1/PAK2/PAK3.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	p21-activated kinase 1;PAK-1;p65-PAK;Alpha-PAK
Accession No.	Swiss-Prot#:Q13153//Q13177//O75914NCBI Gene ID:5058 5062 5063NCBI mRNA#:NM_001128620.1 NM_002577.4 NM_001128166.1 NCBI Protein#: NP_001122092.1 NP_002568.2 NP_001121638.1
Uniprot	Q13153
GeneID	5058;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	61-67 (PAK2) 68-74 (PAK1/3)
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.

Background

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2.

Note: This product is for in vitro research use only