Shc1 (Ab-349) Conjugated Antibody

Catalog No: #C21316

SAB Signalway Antibody

Package Size: #C21316-AF350 100ul #C21316-AF405 100ul #C21316-AF488 100ul

#C21316-AF555 100ul #C21316-AF594 100ul #C21316-AF647 100ul

#C21316-AF680 100ul #C21316-AF750 100ul #C21316-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

2 ccciiption	
Product Name	Shc1 (Ab-349) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total Shc1 protein.
Immunogen Description	Peptide sequence around aa.347~351 (H-Q-Y-Y-N) derived from Human Shc1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	SH2 domain protein C1;SHC;SHC-transforming protein 1;SHCA;Src homology 2
	domain-containing-transforming protein C1
Accession No.	Swiss-Prot#:P29353NCBI Gene ID:6464NCBI mRNA#:NM_001130040.1 NCBI Protein#:NP_001123512.1
Uniprot	P29353
GeneID	6464;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	46 52 66
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.

Background

Signaling adapter that couples activated growth factor receptors to signaling pathway. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span

Note: This product is for in vitro research use only