## DOK4 Conjugated Antibody

Catalog No: #C32025



 Package Size:
 #C32025-AF350 100ul
 #C32025-AF405 100ul
 #C32025-AF488 100ul

 #C32025-AF555 100ul
 #C32025-AF594 100ul
 #C32025-AF647 100ul

 #C32025-AF680 100ul
 #C32025-AF750 100ul
 #C32025-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

Product Name	DOK4 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total DOK4 protein.
Immunogen Description	Recombinant Protein of human DOK4.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	DOK4;dockingprotein4;FLJ10488;Insulinreceptorsubstrate5;IRS5
Accession No.	Swiss-Prot#:Q8TEW6NCBI Gene ID:55715
Uniprot	Q8TEW6
GenelD	55715;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	37
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## **Application Details**

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250		
AF405 conjugated: most applications: 1: 50 - 1: 250		
AF488 conjugated: most applications: 1: 50 - 1: 250		
AF555 conjugated: most applications: 1: 50 - 1: 250		
AF594 conjugated: most applications: 1: 50 - 1: 250		
AF647 conjugated: most applications: 1: 50 - 1: 250		
AF680 conjugated: most applications: 1: 50 - 1: 250		
AF750 conjugated: most applications: 1: 50 - 1: 250		
Biotin conjugated: working with enzyme-conjugated str		

Antibodies were purified by affinity purification using immunogen.

## Background

The downstream of kinase family (Dok-1-7) are members of a class of docking proteins that interact with receptor tyrosine kinases and, via this interaction, mediate biological responses within the body. Dok-4 (Downstream of kinase-4) is a 326 amino acid protein that contains one PH domain and one IRS-type PTB domain and belongs to the Dok family of interacting proteins. Expressed in a variety of tissues with highest expression in liver, heart, kidney and skeletal muscle, Dok-4 plays an important role in Ret-mediated neurite outgrowth and may link Ret with downstream effectors during neuronal differentiation. Additionally, Dok-4 is thought to play a positive role in the activation of MAPK pathways and may participate in T-cell induced immune system regulation. Overexpression of Dok-4 is associated with clear cell renal cell carcinoma, suggesting a role for Dok-4 in tumorigenesis.

Note: This product is for in vitro research use only