# NOS1 Conjugated Antibody

Catalog No: #C32287

SAB Signalway Antibody

Package Size: #C32287-AF350 100ul #C32287-AF405 100ul #C32287-AF488 100ul

#C32287-AF555 100ul #C32287-AF594 100ul #C32287-AF647 100ul

#C32287-AF680 100ul #C32287-AF750 100ul #C32287-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

Product Name	NOS1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total NOS1 protein.
Immunogen Description	Recombinant Protein of human NOS1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	NOS1;IHPS1;N-NOS;NC-NOS;NOS
Accession No.	Swiss-Prot#:P29475NCBI Gene ID:4842
Uniprot	P29475
GeneID	4842;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	160
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

### **Application Details**

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

#### **Product Description**

Antibodies were purified by affinity purification using immunogen.

#### Background

Nitric Oxide Synthase (NOS) catalyses the formation of nitric oxide (NO) and citruline from L-arginine, oxygen and cofactors. Three family members have been characterized: neuronal NOS (nNOS), which is found primarily in neuronal tissue; inducible NOS (iNOS), which is induced by interferon gamma and lipopolysaccharides in the kidney and cardiovascular system; and endothelial NOS (eNOS), which is expressed in blood vessels (1). NO is a messenger molecule with diverse functions throughout the body including the maintenance of vascular integrity, homeostasis, synaptic plasticity, long-term potentiation, learning, and memory (2,3).

nNOS binds selectively to the second PDZ domain of PSD-95, and the enzymatic activity of nNOS is either downregulated by phosphorylation at Ser847 by CaM-KII or stimulated by calcium influxes governed by the NMDA receptor channels (4-5).

Note: This product is for in vitro research use only