

IDE Conjugated Antibody

Catalog No: #C32354



Package Size: #C32354-AF350 100ul #C32354-AF405 100ul #C32354-AF488 100ul
 #C32354-AF555 100ul #C32354-AF594 100ul #C32354-AF647 100ul
 #C32354-AF680 100ul #C32354-AF750 100ul #C32354-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | IDE Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous level of total IDE protein. |
| Immunogen Description | Recombinant protein of human IDE. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | FLJ35968;INSULYSIN |
| Accession No. | Swiss-Prot#:P14735NCBI Gene ID:3416 |
| Uniprot | P14735 |
| GeneID | 3416; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 118 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were purified by affinity purification using immunogen.

Background

This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulin's activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheimer's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causative for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described but have not been experimentally verified.

Note: This product is for in vitro research use only