ITGAL Conjugated Antibody

Catalog No: #C32362



 Package Size:
 #C32362-AF350 100ul
 #C32362-AF405 100ul
 #C32362-AF488 100ul

 #C32362-AF555 100ul
 #C32362-AF594 100ul
 #C32362-AF647 100ul

 #C32362-AF680 100ul
 #C32362-AF750 100ul
 #C32362-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	ITGAL Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total ITGAL protein.
Immunogen Description	Recombinant Protein of human ITGAL.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	CD11A;LFA-1;LFA1A
Accession No.	Swiss-Prot#:P20701NCBI Gene ID:3683
Uniprot	P20701
GenelD	3683;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	129
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250		
AF405 conjugated: most applications: 1: 50 - 1: 250		
AF488 conjugated: most applications: 1: 50 - 1: 250		
AF555 conjugated: most applications: 1: 50 - 1: 250		
AF594 conjugated: most applications: 1: 50 - 1: 250		
AF647 conjugated: most applications: 1: 50 - 1: 250		
AF680 conjugated: most applications: 1: 50 - 1: 250		
AF750 conjugated: most applications: 1: 50 - 1: 250		
Biotin conjugated: working with enzyme-conjugated str		

Antibodies were purified by affinity purification using immunogen.

Background

ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene.

Note: This product is for in vitro research use only