

SPIN1 Conjugated Antibody

Catalog No: #C33091



Package Size: #C33091-AF350 100ul #C33091-AF405 100ul #C33091-AF488 100ul
 #C33091-AF555 100ul #C33091-AF594 100ul #C33091-AF647 100ul
 #C33091-AF680 100ul #C33091-AF750 100ul #C33091-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | SPIN1 Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of total SPIN1 protein. |
| Immunogen Description | Recombinant protein of human SPIN1. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | SPIN |
| Accession No. | Swiss-Prot#:Q9Y657NCBI Gene ID:10927 |
| Uniprot | Q9Y657 |
| GeneID | 10927; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 29 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were purified by affinity purification using immunogen.

Background

Chromatin reader that specifically recognizes and binds histone H3 both trimethylated at 'Lys-4' and asymmetrically dimethylated at 'Arg-8' (H3K4me3 and H3R8me2a) and acts as an activator of Wnt signaling pathway downstream of PRMT2. In case of cancer, promotes cell cancer proliferation via activation of the Wnt signaling pathway . Overexpression induces metaphase arrest and chromosomal instability. Localizes to active rDNA loci and promotes the expression of rRNA genes . May play a role in cell-cycle regulation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process that takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulating maternal transcripts to control meiotic resumption.

Note: This product is for in vitro research use only