

AurB/C (Ab-202/175) Conjugated Antibody

Catalog No: #C33177



Package Size: #C33177-AF350 100ul #C33177-AF405 100ul #C33177-AF488 100ul
 #C33177-AF555 100ul #C33177-AF594 100ul #C33177-AF647 100ul
 #C33177-AF680 100ul #C33177-AF750 100ul #C33177-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	AurB/C (Ab-202/175) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total AurB/C protein.
Immunogen Description	Synthesized non-phosphopeptide derived from human AurB/C around the phosphorylation site of threonine 202/175 (C-G-T(p)-L-D).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	AIE1;AIE2;AURKC;Aurora-C;Aurora/Ipl1-related kinase 3
Accession No.	Swiss-Prot#:Q9UQB9NCBI Gene ID:9212/6795
Uniprot	Q9UQB9
GeneID	6795;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	39
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Serine/threonine-protein kinase component of the chromosomal passenger complex (CPC), a complex that acts as a key regulator of mitosis. The CPC complex has essential functions at the centromere in ensuring correct chromosome alignment and segregation and is required for chromatin-induced microtubule stabilization and spindle assembly. Plays also a role in meiosis and more particularly in spermatogenesis. Has redundant cellular functions with AURKB and can rescue an AURKB knockdown. Like AURKB, AURKC phosphorylates histone H3 at 'Ser-10' and 'Ser-28'. Phosphorylates TACC1, another protein involved in cell division, at 'Ser-228'.

Note: This product is for in vitro research use only