

TOB1 (Ab-164) Conjugated Antibody

Catalog No: #C33296



Package Size: #C33296-AF350 100ul #C33296-AF405 100ul #C33296-AF488 100ul
 #C33296-AF555 100ul #C33296-AF594 100ul #C33296-AF647 100ul
 #C33296-AF680 100ul #C33296-AF750 100ul #C33296-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | TOB1 (Ab-164) Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total TOB1 protein. |
| Immunogen Description | Synthesized non-phosphopeptide derived from human TOB1 around the phosphorylation site of serine 164 (A-V-S(p)-P-T). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | TOB; Transducer of erbB-2 1 |
| Accession No. | Swiss-Prot#:P50616NCBI Gene ID:10140 |
| Uniprot | P50616 |
| GeneID | 10140; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 40 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Anti-proliferative protein; the function is mediated by association with deadenylase subunits of the CCR4-NOT complex.

Note: This product is for in vitro research use only