

## GluR7 Conjugated Antibody

Catalog No: #C33390



Package Size: #C33390-AF350 100ul #C33390-AF405 100ul #C33390-AF488 100ul  
 #C33390-AF555 100ul #C33390-AF594 100ul #C33390-AF647 100ul  
 #C33390-AF680 100ul #C33390-AF750 100ul #C33390-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	GluR7 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of total GluR7 protein.
Immunogen Description	Synthesized peptide derived from human GluR7.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	EAA5;GLR7;GluR7a;dJ1090M5.1 (glutamate receptor;ionotropic
Accession No.	Swiss-Prot#:Q14831NCBI Gene ID:2917
Uniprot	Q14831
GeneID	2917;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	100
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Product Description

---

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

## Background

---

G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.

---

Note: This product is for in vitro research use only