

## TACC1 Conjugated Antibody

Catalog No: #C33610



Package Size: #C33610-AF350 100ul #C33610-AF405 100ul #C33610-AF488 100ul  
 #C33610-AF555 100ul #C33610-AF594 100ul #C33610-AF647 100ul  
 #C33610-AF680 100ul #C33610-AF750 100ul #C33610-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	TACC1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous levels of total TACC1 protein.
Immunogen Description	Synthesized peptide derived from N-terminal of human TACC1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	gastric cancer antigen Ga55;KIAA1103;taxin 1;transforming acidic coiled-coil-containing protein 1
Accession No.	Swiss-Prot#:O75410NCBI Gene ID:6867
Uniprot	O75410
GeneID	6867;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	87
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Product Description

---

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

## Background

---

Likely involved in the processes that promote cell division prior to the formation of differentiated tissues.

---

Note: This product is for in vitro research use only