

ZNF638 Conjugated Antibody

Catalog No: #C33886



Package Size: #C33886-AF350 100ul #C33886-AF405 100ul #C33886-AF488 100ul
 #C33886-AF555 100ul #C33886-AF594 100ul #C33886-AF647 100ul
 #C33886-AF680 100ul #C33886-AF750 100ul #C33886-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	ZNF638 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total ZNF638 protein.
Immunogen Description	Synthesized peptide derived from internal of human ZNF638.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Zinc finger protein 638; Nuclear protein 220; Zinc finger matrix-like protein; Cutaneous T-cell lymphoma-associated antigen se33-1; CTCL tumor antigen se33-1
Accession No.	Swiss-Prot#:Q14966NCBI Gene ID:27332
Uniprot	Q14966
GeneID	27332;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	220
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Early regulator of adipogenesis that works as a transcription cofactor of CEBPs, controlling the expression of PPARG and probably of other proadipogenic genes, such as SREBF1 By similarity. Binds to cytidine clusters in double-stranded DNA.

Note: This product is for in vitro research use only