ATF5 Conjugated Antibody

Catalog No: #C34127



 Package Size:
 #C34127-AF350 100ul
 #C34127-AF405 100ul
 #C34127-AF488 100ul

 #C34127-AF555 100ul
 #C34127-AF594 100ul
 #C34127-AF647 100ul

 #C34127-AF680 100ul
 #C34127-AF750 100ul
 #C34127-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | ATF5 Conjugated Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous levels of total ATF5 protein. |
| Immunogen Description | Synthesized peptide derived from internal of human ATF5. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Cyclic AMP-dependent transcription factor ATF-5; Activating transcription factor 5; Transcription factor |
| | ATFx;ATF5;ATFX |
| Accession No. | Swiss-Prot#:Q9Y2D1NCBI Gene ID:22809 |
| Uniprot | Q9Y2D1 |
| GenelD | 22809; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 31 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |
| | |

Application Details

| Suggested Dilution: |
|---|
| AF350 conjugated: most applications: 1: 50 - 1: 250 |
| AF405 conjugated: most applications: 1: 50 - 1: 250 |
| AF488 conjugated: most applications: 1: 50 - 1: 250 |
| AF555 conjugated: most applications: 1: 50 - 1: 250 |
| AF594 conjugated: most applications: 1: 50 - 1: 250 |
| AF647 conjugated: most applications: 1: 50 - 1: 250 |
| AF680 conjugated: most applications: 1: 50 - 1: 250 |
| AF750 conjugated: most applications: 1: 50 - 1: 250 |

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Transcriptional activator which binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters and blocks the differentiation of neuroprogenitor cells into neurons. Its transcriptional activity is enhanced by CCND3 and slightly inhibited by CDK4.

Note: This product is for in vitro research use only