

LFNG Conjugated Antibody

Catalog No: #C34489



Package Size: #C34489-AF350 100ul #C34489-AF405 100ul #C34489-AF488 100ul
 #C34489-AF555 100ul #C34489-AF594 100ul #C34489-AF647 100ul
 #C34489-AF680 100ul #C34489-AF750 100ul #C34489-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	LFNG Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total LFNG protein.
Immunogen Description	Synthesized peptide derived from internal of human LFNG.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	beta-1;3-N-acetylglucosaminyltransferase lunatic fringe;EC 2.4.1.222;LFNG O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase;O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase
Accession No.	Swiss-Prot#:Q8NES3NCBI Gene ID:3955
Uniprot	Q8NES3
GeneID	3955;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	42
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Glycosyltransferase that initiates the elongation of O-linked fucose residues attached to EGF-like repeats in the extracellular domain of Notch molecules. Decreases the binding of JAGGED1 to NOTCH2 but not that of DELTA1. Essential mediator of somite segmentation and patterning By similarity.

Note: This product is for in vitro research use only