

BORG4 Conjugated Antibody

Catalog No: #C34558



Package Size: #C34558-AF350 100ul #C34558-AF405 100ul #C34558-AF488 100ul
 #C34558-AF555 100ul #C34558-AF594 100ul #C34558-AF647 100ul
 #C34558-AF680 100ul #C34558-AF750 100ul #C34558-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	BORG4 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total BORG4 protein.
Immunogen Description	Synthesized peptide derived from internal of human BORG4.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	binder of Rho GTPases 4;BORG4;CDC42 effector protein (Rho GTPase binding) 4;Cdc42 effector protein 4;CEP4
Accession No.	Swiss-Prot#:Q9H3Q1NCBI Gene ID:23580
Uniprot	Q9H3Q1
GeneID	23580;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	36
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. Induces pseudopodia formation, when overexpressed in fibroblasts.

Note: This product is for in vitro research use only