

RNF138 Conjugated Antibody

Catalog No: #C34663



Package Size: #C34663-AF350 100ul #C34663-AF405 100ul #C34663-AF488 100ul
 #C34663-AF555 100ul #C34663-AF594 100ul #C34663-AF647 100ul
 #C34663-AF680 100ul #C34663-AF750 100ul #C34663-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	RNF138 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total RNF138 protein.
Immunogen Description	Synthesized peptide derived from internal of human RNF138.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	E3 ubiquitin-protein ligase RNF138;EC 6.3.2.-;RING finger protein 138;Nemo-like kinase-associated RING finger protein;NLK-associated RING finger protein
Accession No.	Swiss-Prot#:Q8WVD3NCBI Gene ID:51444
Uniprot	Q8WVD3
GeneID	51444;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	28
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

E3 ubiquitin-protein ligase. Together with NLK, involved in the ubiquitination and degradation of TCF/LEF. Also exhibits auto-ubiquitination activity in combination with UBE2K. May act as a negative regulator in the Wnt/beta-catenin-mediated signaling pathway.

Note: This product is for in vitro research use only