

MT-ND1 Conjugated Antibody

Catalog No: #C34839



Package Size: #C34839-AF350 100ul #C34839-AF405 100ul #C34839-AF488 100ul
 #C34839-AF555 100ul #C34839-AF594 100ul #C34839-AF647 100ul
 #C34839-AF680 100ul #C34839-AF750 100ul #C34839-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	MT-ND1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total MT-ND1 protein.
Immunogen Description	Synthesized peptide derived from internal of human MT-ND1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	NADH-ubiquinone oxidoreductase chain 1;EC 1.6.5.3;NADH dehydrogenase subunit 1;MT-ND1;MTND1
Accession No.	Swiss-Prot#:P03886NCBI Gene ID:4535
Uniprot	P03886
GeneID	4535;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	36
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone By similarity. HAMAP-Rule MF_01350

Note: This product is for in vitro research use only