KCNK15 Conjugated Antibody

Catalog No: #C34912

SAB Signalway Antibody

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size: #C34912-AF350 100ul #C34912-AF405 100ul #C34912-AF488 100ul

#C34912-AF555 100ul #C34912-AF594 100ul #C34912-AF647 100ul

#C34912-AF680 100ul #C34912-AF750 100ul #C34912-Biotin 100ul

Description

| Product Name | KCNK15 Conjugated Antibody |
|-----------------------|---------------------------------------------------------------------------------------------|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total KCNK15 protein. |
| Immunogen Description | Synthesized peptide derived from internal of human KCNK15. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Acid-sensitive potassium channel protein TASK-5; Potassium channel subfamily K member |
| | 15;TASK5;TWIK-related acid-sensitive K(+) channel 5;Two pore potassium channel KT3.3 |
| Accession No. | Swiss-Prot#:Q9H427NCBI Gene ID:60598 |
| Uniprot | Q9H427 |
| GeneID | 60598; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 42 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Probable potassium channel subunit. No channel activity observed in heterologous systems. May need to associate with another protein to form a functional channel.

Note: This product is for in vitro research use only