# UBAP2L Conjugated Antibody

Catalog No: #C35131

Package Size: #C35131-AF350 100ul #C35131-AF405 100ul #C35131-AF488 100ul #C35131-AF555 100ul #C35131-AF594 100ul #C35131-AF647 100ul #C35131-AF680 100ul #C35131-AF750 100ul #C35131-Biotin 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

## Description

| | |
|---|---|
| Product Name | UBAP2L Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total UBAP2L protein. |
| Immunogen Description | Synthesized peptide derived from internal of human UBAP2L. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | UBAP2L;ubiquitin-associated protein 2-like;UBP2L |
| Accession No. | Swiss-Prot#:Q14157NCBI Gene ID:9898 |
| Uniprot | Q14157 |
| GeneID | 9898; |
| Excitation Emission | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Calculated MW | 115 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

## Background

a ubiquitous protein containing an UBA-domain, which are present in multiple proteins of the ubiquitination pathway. Encoded by a gene of the human epidermal differentiation complex (EDC). May be involved in epidermal differentiation. Three isoforms of the human protein are produced by alternative splicing.

Note: This product is for in vitro research use only