# **TNFRSF10D** Conjugated Antibody

Catalog No: #C35312



 Package Size:
 #C35312-AF350 100ul
 #C35312-AF405 100ul
 #C35312-AF488 100ul

 #C35312-AF555 100ul
 #C35312-AF594 100ul
 #C35312-AF647 100ul

 #C35312-AF680 100ul
 #C35312-AF750 100ul
 #C35312-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

# Description

Product Name	TNFRSF10D Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Ни
Specificity	The antibody detects endogenous levels of total TNFRSF10D protein.
Immunogen Description	Synthesized peptide derived from internal of human TNFRSF10D.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Tumor necrosis factor receptor superfamily member 10D;Decoy receptor 2;DcR2;TNF-related
	apoptosis-inducing ligand receptor 4;TRAIL receptor 4
Accession No.	Swiss-Prot#:Q9UBN6NCBI Gene ID:8793
Uniprot	Q9UBN6
GeneID	8793;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	42
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

### **Application Details**

Suggested Dilution:
AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

### **Product Description**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

# Background

Receptor for the cytotoxic ligand TRAIL. Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis. Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to it cannot but according to it can induce the NF-kappa-B pathway.

Note: This product is for in vitro research use only