

## DKK4 Conjugated Antibody

Catalog No: #C35713



Package Size: #C35713-AF350 100ul #C35713-AF405 100ul #C35713-AF488 100ul  
 #C35713-AF555 100ul #C35713-AF594 100ul #C35713-AF647 100ul  
 #C35713-AF680 100ul #C35713-AF750 100ul #C35713-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	DKK4 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total DKK4 protein.
Immunogen Description	Fusion protein corresponding to a region derived from internal residues of human dickkopf WNT signaling pathway inhibitor 4
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	DKK-4
Accession No.	Swiss-Prot#:Q9UBT3NCBI Gene ID:27121NCBI Protein#:BC107047
Uniprot	Q9UBT3
GeneID	27121;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. Activity of this protein is modulated by binding to the Wnt co-receptor and the co-factor kremen 2.

---

Note: This product is for in vitro research use only