SELM Conjugated Antibody

Catalog No: #C35911



 Package Size:
 #C35911-AF350 100ul
 #C35911-AF405 100ul
 #C35911-AF488 100ul

 #C35911-AF555 100ul
 #C35911-AF594 100ul
 #C35911-AF647 100ul

 #C35911-AF680 100ul
 #C35911-AF750 100ul
 #C35911-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | SELM Conjugated Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total SELM protein. |
| Immunogen Description | Fusion protein corresponding to residues near the C terminal of human elenoprotein M |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | SEPM |
| Accession No. | Swiss-Prot#:Q8WWX9NCBI Gene ID:140606NCBI Protein#:BC013421 |
| Uniprot | Q8WWX9 |
| GeneID | 140606; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |
| | |

Application Details

| Suggested Dilution: | |
|---|--|
| AF350 conjugated: most applications: 1: 50 - 1: 250 | |
| AF405 conjugated: most applications: 1: 50 - 1: 250 | |
| AF488 conjugated: most applications: 1: 50 - 1: 250 | |
| AF555 conjugated: most applications: 1: 50 - 1: 250 | |
| AF594 conjugated: most applications: 1: 50 - 1: 250 | |
| AF647 conjugated: most applications: 1: 50 - 1: 250 | |
| AF680 conjugated: most applications: 1: 50 - 1: 250 | |
| AF750 conjugated: most applications: 1: 50 - 1: 250 | |
| Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000 | |

Background

This gene encodes a selenoprotein, which contains a selenocysteine (Sec) residue at its active site. The selenocysteine is encoded by the UGA codon that normally signals translation termination. The 3' UTR of selenoprotein genes have a common stem-loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. This gene is expressed in a variety of tissues, and the protein is localized to the perinuclear structures.?

Note: This product is for in vitro research use only