APOA2 Antibody

Catalog No: #45172

Package Size: #45172-1 50ul #45172-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | |
|-------------------|-------|-----|-----|
| | escri | nti | n |
| $\boldsymbol{ u}$ | COUL | μu | ULI |

| Product Name | APOA2 Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | The antibody was purified by immunogen affinity chromatography. |
| Applications | WB, IF/IC |
| Species Reactivity | Hu Ms Rt |
| Specificity | APOA2 Antibody detects endogenous levels of total APOA2 |
| Immunogen Type | Peptide |
| Immunogen Description | KLH-conjugated synthetic peptide of human Apolipoprotein A2 |
| Target Name | APOA2 |
| Other Names | APOA2, Apo-AII, ApoAII, ApoA-II, Apolipoprotein A-II, Apolipoprotein A2, Apolipoprotein A II |
| Accession No. | Swiss-Prot#: P02652NCBI Gene ID:815 |
| Uniprot | P02652 |
| GeneID | 336; |
| Calculated MW | 11 kDa |
| Concentration | 1.0mg/mL |
| Formulation | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium |
| | azide. |
| Storage | Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles. |
| | |


Application Details

WB (1/500 - 1/2000), IF/IC (1/50 - 1/200)

Images



Western blot analysis of Apolipoprotein A2 expression in Human plasma (A) whole cell lysates.



Immunofluorescent analysis of Apolipoprotein A2 staining in C6 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Note: This product is for in vitro research use only