S100A6 Conjugated Antibody

Catalog No: #C37888



Package Size: #C37888-AF350 100ul #C37888-AF405 100ul #C37888-AF488 100ul

#C37888-AF555 100ul #C37888-AF594 100ul #C37888-AF647 100ul

#C37888-AF680 100ul #C37888-AF750 100ul #C37888-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	S100A6 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total S100A6 protein.
Immunogen Description	Synthetic peptide corresponding to residues near the N terminal of human S100 calcium binding protein A6
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	2A9; PRA; 5B10; CABP; CACY
Accession No.	Swiss-Prot#:P06703NCBI Gene ID:6277NCBI Protein#:NP_000531/P21817
Uniprot	P06703
GeneID	6277;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

 $Biotin \ conjugated: working \ with \ enzyme-conjugated \ streptavidin, \ most \ applications: \ 1:50 - 1:1,000$

Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in stimulation of Ca2+-dependent insulin release, stimulation of prolactin secretion, and exocytosis.?

Note: This product is for in vitro research use only