

KLRK1 Conjugated Antibody

Catalog No: #C38721



Package Size: #C38721-AF350 100ul #C38721-AF405 100ul #C38721-AF488 100ul
 #C38721-AF555 100ul #C38721-AF594 100ul #C38721-AF647 100ul
 #C38721-AF680 100ul #C38721-AF750 100ul #C38721-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	KLRK1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous level of total KLRK1 antibody.
Immunogen Description	Recombinant protein of human KLRK1.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	KLR; CD314; NKG2D; NKG2-D; D12S2489E;
Accession No.	Swiss-Prot#:P26718NCBI Gene ID:22914
Uniprot	P26718
GeneID	100528032;22914;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	25
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.

Note: This product is for in vitro research use only