

Integrin-linked protein kinase Polyclonal Conjugated Antibody

Catalog No: #C42192

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42192-AF350 100ul #C42192-AF405 100ul #C42192-AF488 100ul

#C42192-AF555 100ul #C42192-AF594 100ul #C42192-AF647 100ul

#C42192-AF680 100ul #C42192-AF750 100ul #C42192-Biotin 100ul

Description

Product Name	Integrin-linked protein kinase Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Integrin-linked protein kinase polyclonal antibody.
Immunogen Description	Recombinant human Integrin-linked protein kinase protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	59 kDa serine/threonine-protein kinase,ILK-1,ILK-2,p59ILK
Accession No.	Swiss-Prot#:Q13418
Uniprot	Q13418
GeneID	3611;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Receptor-proximal protein kinase regulating integrin-mediated signal transduction. May act as a mediator of inside-out integrin signaling. Focal adhesion protein part of the complex ILK-PINCH. This complex is considered to be one of the convergence points of integrin- and growth factor-signaling pathway. Could be implicated in mediating cell architecture, adhesion to integrin substrates and anchorage-dependent growth in epithelial cells. Phosphorylates beta-1 and beta-3 integrin subunit on serine and threonine residues, but also AKT1 and GSK3B.

Note: This product is for in vitro research use only